

Portland Special Status Species Plants (updated 2009)

| SCIENTIFIC NAME | COMMON NAME | Site conditions and habitats found |
|--|------------------------------------|--|
| <i>Agrostis howellii</i> | Howell's bentgrass | Moist, shady cliffs; canyon walls; waterfall spray zones; talus |
| <i>Cimicifuga elata</i> var. <i>elata</i> | Tall bugbane | In, or at the margins of, moist, mixed coniferous and deciduous forest. Areas of filtered light. |
| <i>Delphinium leucophaeum</i> | White rock larkspur | Mesic woodlands and forests |
| <i>Delphinium pavonaceum</i> | Peacock larkspur | Prairie |
| <i>Howellia aquatilis</i> | Howellia | Ponds or sloughs / submerged or partially flooded |
| <i>Rorippa columbiae</i> | Columbia cress | Near water; seasonally inundated |
| <i>Sericocarpus rigidus</i> (syn <i>Aster curtus</i>) | White-topped aster | Grasslands |
| <i>Sullivantia oregana</i> | Oregon sullivantia | Moist cliffs, especially near waterfalls; growing on small pockets of basalt derived soil. |
| <i>Castilleja levisecta</i> | Golden paintbrush | Seasonally (spring) wet prairies |
| <i>Artemisia campestris</i> var. <i>wormskioldii</i> | Northern wormwood | Rocky, cobbly shorelines and riverbanks |
| <i>Carex comosa</i> | Bristly sedge | Marshes, lake shores, wet meadows |
| <i>Carex retrorsa</i> | Retrorse sedge | Wet thickets, swamps, marshes, lake shores |
| <i>Delphinium nuttallii</i> | Nuttall's larkspur | Low moist ground, gravelly outwashes, basalt cliffs |
| <i>Fritillaria camschatcensis</i> | Indian rice / black lilly | Moist open meadows, near lakes and streams, coniferous-forested wetlands |
| <i>Heliotropium curassavicum</i> | Salt heliotrope | Dry or moist saline and alkaline areas |
| <i>Rotala ramosior</i> | Toothcup | Wet, swampy places; lake and pond margins; along rivers |
| <i>Wolffia columbiana</i> | Columbia water-meal | Freshwater ponds, lakes, and slow-moving streams |
| <i>Sedella pumila</i> | Sierra mock-stonecrop | Thin soil on rock outcrops, gravelly soil, vernal pools. |
| <i>Ammannia robusta</i> | Grand redstem (loosestrife family) | Riverine emergent wetlands, riparian mudflat wetlands |
| <i>Elodea nuttallii</i> | Nuttall's waterweed | In lakes, ponds, ditches |
| <i>Hierochloe odorata</i> | Holy grass | Moist slopes, meadows, stream banks |

| | | |
|------------------------------|--------------------------|--|
| <i>Polygonum punctatum</i> | Dotted smartweed | Swamps, shallows; pond, stream, and lake edges |
| <i>Scirpus pallidus</i> | Pale bulrush | Wet meadows, stream banks, springs |
| <i>Zizia aptera</i> | Golden alexanders | Prairie, rocky upland woodlands, limestone glades |
| <i>Bergia texana</i> | Texas bergia | Sand bars, vernal pools, moist disturbed soil |
| <i>Bolandra oregana</i> | Oregon bolandra | Wooded areas on cliffs near waterfalls |
| <i>Cypripedium montanum</i> | Mountain lady's-slipper | Mid- to late-successional forest communities; mixed conifers or mixed evergreen/oak woodland |
| <i>Euonymus occidentalis</i> | | Moist, wooded/forested areas |
| <i>Montia howellii</i> | Howell's montia | Moist, disturbed soils; lowlands |
| <i>Poa laxiflora</i> | Loose-flowered bluegrass | Open meadows along stream banks; upper margins of sea beaches |
| <i>Poa marcida</i> | Weak bluegrass | |
| <i>Sidalcea campestris</i> | Meadow checker-mallow | Prairie / grasslands (roadside ditches) |