



Vegetation Unit Summaries for **Overlook House Community Center (OVHS)**

Report date: 7/29/2009

'\*' = non-native invasive species

Visit data as of: 7/29/2009

**Unit OVHS\*001**

**Size:** 0.99 Acres

**NVCS Class:** Forest

**NVCS Subclass:** Mixed evergreen-deciduous forest

**NVCS Group:** Mixed needle-leaved evergreen - cold-deciduous forest

**NVCS SubGroup:** Natural / Semi-natural

**NVCS Ecological System:** unknown

**Landform:** Side-hill, Upper 3rd

**Slope:** Steep (20 to 30%)

**Aspect:** Southwest

**Notes:**

**Visit date:** 10/4/2006

**Ecological Health:** Poor.

**% Tree canopy:** 55%

**% Non-Native Cover:** 65

**General Note:**

**Management Note:**

**Eco Note:**

**Primary Management concerns:** Invasives, Erosion, Trampling, Litter, Yard Debris, Informal Trails.

**Visit Species:**

	COVER CLASS	DOMINANT	DBH	REGENERATING	PLANTED
Rubus discolor (Himalayan blackberry)	20% to 50%	Y			
Hedera helix (English ivy)	20% to 50%				
Thuja plicata (western red cedar)	10% to 20%	Y	10-20"		
Acer macrophyllum (bigleaf maple)	10% to 20%		10-20"		
Clematis vitalba (clematis)	10% to 20%				
Corylus cornuta (western beaked hazelnut)	10% to 20%				
Sequoiadendron giganteum (giant redwood)	1% to 10%	Y	>30"		
Carya ovata (shagbark hickory)	1% to 10%		10-20"		
Juglans nigra (black walnut)	1% to 10%		5-10"		
Populus balsamifera ssp. trichocarpa (black cott)	1% to 10%		5-10"	Y	
Prunus avium (sweet pie cherry)	1% to 10%		5-10"		
Conium maculatum (poison hemlock)	1% to 10%				
Convolvulus arvensis (morning glory)	1% to 10%				
Ilex aquafolium (English holly)	1% to 10%				
Solanum dulcamara (bittersweet nightshade)	1% to 10%				
Pinus contorta (lodgepole pine)	trace		0-5"	Y	
Quercus garryana (Oregon white oak)	trace		0-5"	Y	
Sorbus aucuparia (European mountain ash)	trace		0-5"		
Acer circinatum (vine maple)	trace				
Crataegus monogyna (European hawthorn)	trace				
Cytisus scoparius (Scot's broom)	trace				
Daucus carota (Queen Anne's lace)	trace				
Epilobium ciliatum (willowherb)	trace				
Gaultheria shallon (salal)	trace				
Hypericum perforatum (common St. John's wort)	trace				

Lactuca muralis (wall lettuce)	trace
Philadelphus lewisii (Lewis' mockorange)	trace
Polystichum munitum (sword fern)	trace
Pteridium aquilinum (bracken fern)	trace
Rhododendron sp. (rhododendron)	trace
Ribes sanguineum (red-flowering currant)	trace
Rubus parviflorus (thimbleberry)	trace
Rubus ursinus (trailing blackberry)	trace
Sonchus asper (spiny sowthistle)	trace
Streptopus amplexifolius (clasping-leaf twisted-	trace
Buddleia davidii (butterfly bush)	trace

## Unit OVHS\*002

**Size:** 0.38 Acres

**NVCS Class:** Herbaceous

**NVCS Subclass:** Perennial graminoid vegetation

**NVCS Group:** Temperate grassland with a sparse tree layer

**NVCS SubGroup:** Cultural

**NVCS Ecological System:** unknown

**Landform:** unknown

**Slope:** unknown

**Aspect:** unknown

**Visit date:** 10/4/2006

**Ecological Health:** Poor.

**% Tree canopy:** 5%

**% Non-Native Cover:** 80

**General Note:**

Landscaped gardens and building in this site. grass is 35%.

**Management Note:**

**Eco Note:**

**Visit Species:**

	COVER CLASS	DOMINANT	DBH	REGENERATING	PLANTED
Rhododendron sp. (rhododendron)	10% to 20%				
Ilex aquafolium (English holly)	1% to 10%				
Rosa (rose)	1% to 10%				
Calluna (heather)	trace				
Hosta (plantain lily)				trace	